Table 3.—Solar-radiation measurements, and determinations atmospheric turbidity factor, \$\beta\$, Washington, D.C., March 1933

Date and solar hour angle	Solar alti- tude,	Air mass, m	I - cal.	I, cal.	I, cal.	β	Blue- ness of sky	Atmospheric dust particles per cubic centimeter	Notes: Sky light polari zation, P, clouds, etc.
1933 Mar. 6 3:54 a	20-01 21-03 21-45 22-26 31-16 31-20	2. 90 2. 77 2. 69 2. 61 1. 92 1. 92	1. 225 1. 244 1. 269 1. 280 1. 285 1. 257	0. 877 . 928 . 934 . 936 . 880 . 880	0.788 .744 .756 .759 .735	0. 040 . 040 . 038 . 035 . 060 . 070	5	914	P=51.2% Cirri.
Mar. 9 4:18 a	16-34 17-34 24-10 25-10 32-54 33-29 42-08 42-47	3. 48 3. 28 2. 43 2. 34 1. 84 1. 81 1. 49 1. 47	1. 079 1. 105 1. 217 1. 250 1. 340 1. 360 1. 444 1. 463	.791 .797 .881 .879 .918 .921 .921	.665 .667 .707 .718 .746 .747 .735	. 042 . 040 . 045 . 040 . 045 . 045 . 040 . 035	5	275	P=48.2%
Mar. 27 4:51 a	15-00 15-36 19-40 20-35 26-34 27-18	3. 82 3. 68 2. 95 2. 82 2. 23 2. 17	. 645 . 644 . 753 . 787 . 816 . 832	. 488 . 491 . 569 . 571 . 635 . 638	. 447 . 450 . 516 . 521 . 537 . 540	. 130 . 135 . 165 . 165 . 180 . 180	4	899	Cirri. P=42.6%
Mar. 29 4:57 a	14-24 14-58 19-02 19-56 21-36 22-12 26-51 27-35 43-22 43-56 52-20 52-40 48-19	3. 98 3. 84 3. 05 2. 92 2. 70 2. 63 2. 21 1. 46 1. 44 1. 26 1. 34	. 909 . 922 1. 029 1. 058 1. 075 1. 106 1. 211 1. 236 1. 353 1. 338 1. 424 1. 429 1. 350	. 688 . 691 . 765 . 769 . 794 . 853 . 856 . 888 . 891 . 935 . 938 . 938	.591 .596 .647 .650 .671 .672 .700 .700 .700 .759 .759	.065 .065 .075 .065 .078 .072 .060 .052 .105 .115 .068 .068	6	317	P = 53.7 %
1:36 p. 1:42 p. 2:43 p. 2:47 p. 3:26 p. 3:30 p.	48-19 47-36 38-42 37-40 31-14 30-42	1. 34 1. 36 1. 60 1. 63 1. 93 1. 95	1. 306 1. 295 1. 301 1. 218 1. 221	. 950 . 953 . 884 . 887 . 860 . 863	.771 .694 .694 .694 .697	. 140 . 140 . 140 . 140 . 135 . 135			

POSITIONS AND AREAS OF SUN SPOTS

[Communicated by Capt. J. F. Hellweg, Superintendent United States Naval Observatory. Data furnished by Naval Observatory, in cooperation with Harvard, Perkins, and Mount Wilson Observatories. The differences of latitude are measured from central meridian, positive west. The north latitude are plus. Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere. The total area, including spots and groups, is given for each day in the last column]

Date		Eastern standard civil time		Heliographic			Area	
				Longi- tude	Lati- tude	Spot	Group	for each day
1933	h.	m.			•			
Mar. 1 (Naval Observatory)	lii	56	-64.0	305.4	+13.0		123	123
Mar. 2 (Mount Wilson)	12	0	-51.0	305.2	± 14.0		223	
11102. 2 (11204110 1120-)	i		-23.0	333. 2	+9.0	11		234
Mar. 3 (Naval Observatory)	11	34	-37.0	306.3	+13.0		93	93
Mar. 4 (Naval Observatory)	11	35	-24.0	306.1	+13.0		93	93
Mar. 5 (Perkins Observatory)	16	0	-13.5	301.0	+13.0		90	
,			-4.5	310.0	+16.5		90	180
Mar. 6 (Naval Observatory)	11	9	+3.0		+13.0		93	93
Mar. 7 (Mount Wilson)	12	35	+16.0	306.0	+14.0		118	118

POSITIONS AND AREAS OF SUN SPOTS-Continued

Date		Eastern standard civil time		Heliographic			Area	
				Longi- tude	Lati- tude	Spot	Group	for each day
Mar. 8 (Naval Observatory)	h. 11	m. 51	+24.0	° 301. 2	ة +11.0		9	
Mar. 9 (Naval Observatory)	10	5	+33. 0 +37. 0	310. 2	+15. 0 +11. 0	77		8
Mar. 10 (Naval Observatory)	11	8	+46. 0 +60. 0	311.0	$+15.0 \\ +15.0$	77 77		8i 7
Mar. 11 (Naval Observatory)	11	49	+74.0	311.7	+15.0	77		7
Mar. 12 (Perkins Observatory) Mar. 13 (Mount Wilson)	14 11	0 30		No spot No spot		-	<i>-</i>	
Mar. 14 (Mount Wilson)	12	25	1	No spot	s			
Mar. 15 (Naval Observatory) Mar. 16 (Naval Observatory)	10 11	58 35		No spot No spot			-	
Mar. 17 (Naval Observatory)	11	44]	No spot	S			
Mar. 18 (Mount Wilson) Mar. 19 (Mount Wilson)		0 50		No spot No spot				
Mar. 20 (Mount Wilson)	10	45)	No spot	s			
Mar. 21 (Mount Wilson)	11 13	0 28	-73. 0	No spot: 	s +4.0		401	40
Mar. 23 (Mount Wilson)	11	45	-58.0	21.5	+5.0		408	40
Mar. 24 (Naval Observatory)	13 11	44 30	-45.0 -32.0	20.3 21.3	$+4.0 \\ +5.0$	- -	216 234	21
Mar. 25 (Mount Wilson)	111	30	+20.0	73. 3	+10.0		31	26
Mar. 26 (Naval Observatory)	12	44	-18.0 +35.0	21. 5 74. 5	+4.0		247 77	32
Mar. 27 (Mount Wilson)	11	15	-5.0	22. 1	+5.0		200	
Mar. 28 (Naval Observatory)	11	20	+49.0 +8.0	76. 1 21. 9	+10.0 +5.0		185	24
Mar. 29 (Naval Observatory)	10	54	$+62.0 \\ +22.0$	75. 9 22. 9	+10.0 +5.0		93 170	27 17
Mar. 30 (Naval Observatory)		13 10	+36. 0 +49. 0	23. 5 23. 3	+5.0 +4.0		123	12
Mar. 31 (Mount Wilson)		10	7.48.0	20.0	74.0		131	
Mean daily area for March.						[-		12

[Communicated by Capt. J. F. Hellweg, Superintendent United States Naval Observatory. Data furnished by Naval Observatory, in cooperation with Harvard, Perkins, and Mount Wilson Observatories. The differences of longitude are measured from central meridian, positive west. The north latitudes are plus. Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere. The total area, including spots and groups, is given for each day in the last column]

PROVISIONAL SUN-SPOT RELATIVE NUMBERS FOR MARCH 1933

[Dependent alone on observations at Zurich and its station at Arosa] [Data furnished through the courtesy of Prof. W. Brunner, University of Zurich, Switzerland]

March 1933	Relative numbers	March 1933	Relative numbers	March 1933	Relative numbers
1 2 3 4 5	10 10 10 13 11	11 12 13 14 15	7 7 0 0	21	$0 \\ d12 \\ 14 \\ 14 \\ Mc24$
6 7 8 9 10	a11 11 9 8	16 17 18 19 20	0 0 0 0 0	26	29 26 a28 22 d12

Mean: 30 days = 10.0.

a= Passage of an average-sized group through the central meridian. b= Passage of a large group or spot through the central meridian. c= New formation of a center of activity: E, on the eastern part of the sun's disk; W, on the western part; M, in the central zone. d= Entrance of a large or average-sized center of activity on the east limb.

AEROLOGICAL OBSERVATIONS

[Aerological Division, W. R. Gregg, in charge]

By L. T. SAMUELS

Free-air temperatures during March were above normal at all stations except Chicago and Cleveland. (See table 1.) The departures were only of moderate magnitude. Free-air relative humidities averaged above nor-

mal with departures of small to moderate magnitude in most cases.

There were no important deviations in the monthly resultant winds as compared to the normals.